Cotton Varieties Planted 2012 Crop

U. S. DEPARTMENT OF AGRICULTURE
Agricultural Marketing Service - Cotton Program
Memphis, Tennessee
September 5, 2012

mp_cn833

Cotton Varieties Planted, United States
2012 Crop

The Deltapine brand of Upland cottonseed was the most popular planted in the United States for the 2012-2013 season, according to the USDA, Agricultural Marketing Service's Cotton and Tobacco Programs. The Bayer CropScience Fibermax brand was the second most popular followed by Phytogen, Americot, Bayer CropScience Stoneville, All-Tex, and Dyna-Gro.

Transgenic varieties - genetically engineered varieties resistant to worms, herbicides, or both - accounted for about 99 percent of the Upland cotton planted in the United States in 2012. Usage of transgenic varieties in 2012 varied from a high of 100 percent in Florida, Louisiana, and Mississippi, to a low of 93 percent in California. Other states planted from 95-99 percent transgenic.

Deltapine brand varieties were the most popular planted in 2012, accounting for 28.2 percent of the United States acreage. This brand accounted for 42.0 percent of the acreage planted in the southeastern states (Alabama, Florida, Georgia, North Carolina, South Carolina, and Virginia). It accounted for about 34.0 percent in the south central states (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee), 20.5 percent in the southwestern states (Texas, Oklahoma, and Kansas), and 33.4 percent in the western states (Arizona, California, and New Mexico). Deltapine's most popular varieties were DP 1044 B2RF, DP 0912 B2RF, DP 1050 B2RF, and DP 1048 B2RF, accounting respectively for 7.2, 4.3, 3.8, and 2.9 percent of the U.S. Upland cotton acreage.

Bayer CropScience Fibermax brand varieties were the second most popular planted in 2012, accounting for 24.5 percent of the United States acreage. These varieties accounted for 5.2 percent of the acreage planted in the southeastern states, 3.3 percent in the south central states, 39.1 percent in the southwestern states, and 19.8 percent in the western states. The most popular Bayer CropScience Fibermax varieties were FM 9170 B2F, FM 1740 B2F, FM 2484 B2F, and FM 9063 B2F, accounting respectively for about 4.7, 4.1, 3.5, and 1.9 percent of the United States acreage planted to Upland cotton.

Phytogen brand varieties were the third most popular planted in 2012, accounting for 18.4 percent of the United States acreage. They accounted for 41.8 percent of the acreage planted in the southeastern states, 19.4 percent of the acreage in the south central states, 8.0 percent in the southwestern states and 33.7 percent in the western states. The most popular Phytogen brand varieties were PHY 499 WRF, PHY 375 WRF, and PHY 367 WRF, accounting respectively for 9.7, 5.4, and 1.7 percent of the United States acreage planted to Upland cotton.

Americot brand varieties were the fourth most popular planted in 2012. These varieties accounted for about 11.2 percent of the acreage planted. They accounted for 1.3 percent of the acreage planted in the southeastern states, 3.2 percent of the acreage in the south central states, and 18.3 percent in the southwestern states. The most popular Americot varieties were NG 4012 B2RF, NG 4111 RF, and NG 3348 B2RF, accounting respectively for 3.0, 1.8, and 1.6 percent of the United States acreage planted to Upland cotton.

Bayer CropScience Stoneville brand varieties were the fifth most popular and accounted for about 10.0 percent of the U.S. acreage planted in 2012. All-Tex varieties were the sixth most popular and accounted for about 4.7 percent of the 2012 cotton acreage. Dyna-Gro varieties were the seventh most popular and accounted for about 2.4 percent of the 2012 cotton acreage.

Phytogen was the most popular brand of American Pima varieties planted in 2012. Phytogen variety PHY 805 RF accounted for 48.8 percent of the United States Pima acreage and was the most popular variety planted in California (52.3 percent of California Pima acreage). Phytogen's PHY 802 RF was the second most planted American Pima variety and accounted for 12.7 percent of the U.S. crop. Deltapine's DP 340 was the next most popular variety and accounted for 12.3 percent of the U.S. Pima acreage.

Bayer CropScience FM 958 and AFD 2485 were the predominate varieties planted by Organic cotton producers. Other varieties planted by organic producers include All-Tex 7A21, All-Tex LA122, All-Tex A102, Bayer CropScience FM 989, and Seed Source Genetics CT 210.B15

Estimates of the percentage of the various varieties of cotton planted in the United States for 2012 were based on informal surveys made by the Cotton+B9 Program Classing Offices. Those surveyed included ginners, seed dealers, extension agents, and other knowledgeable sources.

Estimated percentage of Upland cotton planted to leading specified varieties, by growth area, 2012 crop

Variety	US	Southeast	South Central	Southwest	Far West
PHY 499 WRF Phytogen	**9.65%**	**29.85%**	**5.58%**	**3.24%**	**5.36%**
DP 1044 B2RF Deltapine	**7.16%**	**0.06%**	**0.03%**	**12.24%**	**7.40%**
PHY 375 WRF Phytogen	**5.44%**	**8.07%**	**12.19%**	**2.20%**	**6.00%**
FM 9170 B2F Bayer CropScience - FiberMax	**4.70%**	**-**	**-**	**8.33%**	**0.21%**
DP 0912 B2RF Deltapine	**4.28%**	**2.61%**	**16.89%**	**0.91%**	**4.12%**
FM 1740 B2F Bayer CropScience - FiberMax	**4.11%**	**1.00%**	**0.36%**	**6.58%**	**3.35%**
ST 5458 B2RF Bayer CropScience - Stoneville	**3.86%**	**1.30%**	**13.89%**	**1.74%**	**2.39%**
DP 1050 B2RF Deltapine	**3.75%**	**15.98%**	**0.15%**	**0.20%**	**1.67%**
FM 2484 B2F Bayer CropScience - FiberMax	**3.48%**	**-**	**-**	**6.05%**	**2.14%**
Epic RF All-Tex	**3.17%**	**-**	**-**	**5.63%**	**-**
NG 4012 B2RF Americot	**3.01%**	**-**	**-**	**5.34%**	**-**
DP 1048 B2RF Deltapine	**2.92%**	**10.52%**	**1.39%**	**0.46%**	**1.99%**
ST 4145 LLB2 Bayer CropScience - Stoneville	**2.74%**	**4.15%**	**10.03%**	**-**	**0.23%**
ST 5288 B2F Bayer CropScience - Stoneville	**2.11%**	**0.92%**	**10.35%**	**-**	**0.88%**
DG 2570 B2RF Dyna-Gro	**1.96%**	**2.00%**	**1.55%**	**2.19%**	**-**
FM 9063 B2F Bayer CropScience - FiberMax	**1.94%**	**-**	**0.03%**	**3.40%**	**0.54%**
NG 4111 RF Americot	**1.77%**	**-**	**-**	**3.14%**	**-**
PHY 367 WRF Phytogen	**1.71%**	**0.99%**	**0.87%**	**2.14%**	**3.76%**
NG 3348 B2RF Americot	**1.63%**	**-**	**-**	**2.89%**	**-**
FM 9180 B2F Bayer CropScience - FiberMax	**1.45%**	**-**	**-**	**2.53%**	**0.68%**

Estimated percentage of Upland cotton planted to leading specified brands by growth area, 2012 crop

Brand	US	Southeast	South Central	Southwest	Far West
Deltapine	28.15%	41.96%	34.04%	20.50%	33.44%
Bayer CropScience - FiberMax	24.45%	5.23%	3.29%	39.09%	19.80%
Phytogen	18.40%	41.84%	19.37%	7.95%	33.67%
Americot	11.19%	1.29%	3.21%	18.31%	-
Bayer CropScience - Stoneville	10.00%	7.31%	37.66%	2.28%	8.66%
All-Tex	4.66%	-	-	8.26%	-
Dyna-Gro	2.39%	2.08%	1.57%	2.92%	-
Miscellaneous	0.40%	0.29%	0.05%	0.32%	4.43%
Croplan Genetics	0.18%	-	0.14%	0.28%	-
Bronco	0.12%	-	0.67%	-	-
Concho	0.05%	-	-	0.09%	-
Seed Source Genetics	*	-	0.01%	-	-

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2012

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
All-Tex																		
Apex B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.79	-	-	0.17	-	0.11
Arid B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.15	-	0.08
Atlas	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-	0.01
Epic RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5.89	-	3.17
Top-Pick	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.14	-	0.07
Patriot + RF	-	-	-	-	-	-	-	-	-	-	-	-	0.82	-	-	1.32	-	0.73
Rapid B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.02	-	0.01
7A21	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.06	-	0.03
A102	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03	-	0.02
Dinero B2RF (81220)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.11	-	0.06
Edge B2RF (81227)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.22	-	0.12
LA122	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.06	-	0.03
Nitro-44 B2RF (81144)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.39	-	0.21
All-Tex Total	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**1.60**	**-**	**-**	**8.57**	**-**	**4.66**
Americot																		
AM 1511 B2RF	0.05	2.97	-	-	-	0.96	1.50	-	0.36	-	0.26	-	-	3.16	0.59	0.73	1.82	0.78
AM 1532 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	2.03	-	-	2.53	-	1.41
AM 1550 B2RF	-	7.49	-	-	-	-	-	-	0.71	-	0.98	-	-	0.31	1.64	0.02	3.75	0.50
NG 1551 RF	-	-	-	-	-	-	15.00	-	-	-	-	-	-	-	-	0.17	-	0.16
NG 2549 B2RF	-	-	-	-	-	-	1.00	-	-	-	0.04	-	0.53	-	-	0.20	-	0.13
NG 3348 B2RF	-	-	-	-	-	-	2.50	-	-	-	-	-	1.14	-	-	2.96	-	1.63
NG 3410 RF	-	-	-	-	-	-	-	-	-	-	-	-	0.22	-	-	1.74	-	0.94
NG 4010 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.44	-	-	0.82	-	0.45
NG 4012 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.61	-	-	5.56	-	3.01
NG 4111 RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3.28	-	1.77
NG 1572 RF	-	-	-	-	-	-	17.50	-	-	-	-	-	-	-	-	0.55	-	0.37
NG 2051 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.08	-	0.05
Americot Total	**0.05**	**10.47**	**-**	**-**	**-**	**0.96**	**37.50**	**-**	**1.07**	**-**	**1.28**	**-**	**4.97**	**3.47**	**2.22**	**18.64**	**5.56**	**11.19**
Bayer CropScience - FiberMax																		
FM 1740 B2F	0.84	0.81	6.20	1.12	-	1.52	7.50	-	0.86	-	0.18	-	15.42	0.13	0.10	6.20	0.53	4.11
FM 1773 LLB2	-	1.05	-	-	-	0.38	-	-	-	-	0.02	-	-	0.05	-	-	0.05	0.10
FM 1845 LLB2	0.33	1.45	-	-	-	4.63	-	-	-	1.03	0.58	-	-	0.41	-	1.24	0.05	1.39
FM 1880 B2F	-	-	-	0.03	-	-	-	-	-	0.12	-	-	-	-	-	0.03	-	0.02
FM 832 LL	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.07	-	0.04
FM 835 LLB2	-	-	-	0.18	-	-	-	-	-	-	-	-	-	-	-	0.42	-	0.23
FM 840 B2F	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.60	-	0.32
FM 9058 F	-	-	0.92	-	-	-	0.50	-	-	-	-	0.56	-	-	-	1.04	-	0.58
FM 9063 B2F	-	-	1.14	-	-	-	-	-	0.15	-	-	-	0.77	-	0.04	3.52	-	1.94
FM 9101 GT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.35	-	0.19
FM 9103 GT	-	-	-	-	-	-	3.50	-	-	-	-	-	-	-	-	1.04	-	0.57
FM 9160 B2F	-	-	-	-	-	-	5.00	-	-	-	-	-	-	-	-	1.61	-	0.89
FM 9170 B2F	-	-	0.45	-	-	-	-	-	-	-	-	-	0.32	-	-	8.70	-	4.70

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2012

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
FM 9180 B2F	-	-	1.07	-	-	-	2.00	-	-	-	-	1.40	0.44	-	-	2.61	-	1.45
FM 958	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.10	-	0.05
FM 958 LL	-	-	-	-	-	-	-	-	-	-	0.01	-	-	0.02	-	1.56	-	0.84
Acala Daytona RF	-	-	-	24.29	-	-	-	-	-	-	-	-	0.23	-	-	-	-	0.29
FM 832	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.24	-	0.13
FM 1944 GLB2	0.03	2.62	0.08	2.49	-	1.02	-	0.07	1.02	2.05	2.52	-	0.55	1.51	3.60	0.31	2.96	0.85
FM 2011 GT	-	-	0.02	2.23	-	-	4.00	-	-	-	-	-	-	-	-	2.11	-	1.18
FM 2484 B2F	-	-	0.04	-	-	-	-	-	-	-	-	13.74	0.33	-	-	6.34	-	3.48
FM 2989 GLB2	-	-	0.38	-	-	-	-	-	-	-	-	8.51	0.12	-	-	1.30	-	0.74
FM 9250 GL	-	-	-	0.51	-	-	4.50	-	-	-	-	-	-	-	-	0.34	-	0.21
FM 8270 GLB2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.24	-	0.13
Bayer CropScience - FiberMax Total	**1.20**	**5.93**	**10.30**	**30.84**	**-**	**7.56**	**27.00**	**0.07**	**2.03**	**3.20**	**3.31**	**24.20**	**18.16**	**2.12**	**3.74**	**39.99**	**3.59**	**24.45**
Bayer CropScience - Stoneville																		
ST 4288 B2F	2.78	0.09	-	0.28	-	0.08	3.00	-	2.52	0.94	1.06	-	2.31	0.14	0.50	0.08	0.73	0.40
ST 4498 B2RF	0.23	0.36	10.47	-	-	-	-	-	2.50	-	0.26	-	-	0.03	0.08	0.13	0.51	0.35
ST 5288 B2F	1.05	4.94	1.66	0.25	-	-	-	51.24	1.66	13.68	4.03	-	-	0.19	-	-	3.79	2.11
ST 5458 B2RF	1.78	17.05	5.02	-	-	1.83	-	0.71	25.23	20.75	0.30	-	-	0.26	0.19	1.82	-	3.86
ST 4145 LLB2	-	6.79	-	0.64	-	1.14	-	-	12.13	5.51	13.68	-	-	5.35	29.47	-	10.22	2.74
ST 5445 LLB2	-	3.85	-	0.22	3.79	0.62	-	0.12	0.30	1.91	0.17	-	-	0.60	0.39	0.23	0.12	0.54
Bayer CropScience - Stoneville Total	**5.83**	**33.09**	**17.15**	**1.39**	**3.79**	**3.68**	**3.00**	**52.07**	**44.34**	**42.78**	**19.49**	**-**	**2.31**	**6.56**	**30.63**	**2.26**	**15.36**	**10.00**
Bronco																		
141	-	2.65	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.12
Bronco Total	**-**	**2.65**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**0.12**
Concho																		
257	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.09	-	0.05
Concho Total	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**0.09**	**-**	**0.05**
Croplan Genetics																		
CG 3220 B2RF	-	0.06	-	-	-	-	-	-	0.10	-	-	-	-	-	0.19	0.05	-	0.04
CG 3520 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.05	-	0.03
CG 3035 RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*	-	*
CG 3787 B2RF	-	-	-	-	-	-	-	0.81	-	-	-	-	-	-	-	0.19	-	0.12
Croplan Genetics Total	**-**	**0.06**	**-**	**-**	**-**	**-**	**-**	**0.81**	**0.10**	**-**	**-**	**-**	**-**	**-**	**0.19**	**0.29**	**-**	**0.18**
Deltapine																		
DP 0912 B2RF	7.38	25.12	8.66	-	-	2.97	17.00	3.46	31.14	7.72	3.08	-	4.05	2.90	11.38	0.65	1.46	4.28
DP 0920 B2RF	1.10	0.14	0.52	-	-	-	-	-	2.31	-	0.19	-	2.62	0.26	1.58	0.28	0.43	0.39
DP 0924 B2RF	0.78	0.53	1.23	0.26	-	-	5.50	-	0.40	-	1.10	-	1.57	0.31	-	0.33	0.20	0.37
DP 0935 B2RF	0.34	0.46	1.03	-	-	0.11	-	-	-	-	0.08	-	4.39	0.56	-	0.90	-	0.66
DP 0949 B2RF	1.06	-	-	-	7.59	0.83	-	0.32	-	-	0.01	-	-	0.76	0.22	*	-	0.23
DP 1028 B2RF	-	0.96	0.54	-	-	-	-	-	0.09	4.08	8.94	-	-	0.88	-	-	6.20	0.69
DP 1032 B2RF	-	-	-	-	-	-	-	-	-	0.77	0.08	-	2.50	-	-	0.45	0.90	0.34
DP 1034 B2RF	2.49	1.68	0.08	-	-	0.48	-	0.98	1.02	13.33	6.04	-	-	1.50	0.32	-	1.90	1.19
DP 104 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	3.15	-	-	0.26	-	0.21
DP 1044 B2RF	0.42	-	13.74	-	-	-	-	0.32	-	-	-	5.22	34.43	0.07	-	11.42	-	7.16

Table 1. Estimated percentage of upland cotton planted to specified varieties, by states and United States, crop of 2012

	AL	AR	AZ	CA	FL	GA	KS	LA	MO	MS	NC	NM	OK	SC	TN	TX	VA	US
DP 1048 B2RF	15.63	0.14	3.76	0.56	12.06	13.16	-	4.01	-	3.74	0.62	-	0.30	5.94	-	0.47	-	2.92
DP 1050 B2RF	18.69	-	2.84	0.89	26.21	21.58	-	0.51	-	0.40	0.35	-	-	4.99	-	0.21	1.35	3.75
DP 1133 B2RF	3.73	4.99	0.14	0.81	-	0.57	-	14.43	-	7.49	0.52	-	-	0.20	-	0.01	-	1.07
DP 1137 B2RF	7.56	-	-	-	3.79	2.42	-	0.99	-	5.22	7.79	-	-	8.98	-	0.11	1.83	1.43
DP 121 RF	-	-	-	-	-	-	-	-	0.69	-	-	-	0.22	-	0.19	-	-	0.03
DP 164 B2RF	-	-	23.24	11.28	-	-	-	-	-	-	-	-	-	-	-	0.17	-	0.60
DP 174 RF	0.30	-	0.23	0.26	-	-	-	-	0.04	1.05	*	-	0.44	-	-	0.60	-	0.40
DP 1212 B2RF	0.04	-	0.45	-	-	-	1.50	-	-	-	0.06	-	0.22	0.05	-	0.45	0.29	0.27
DP 1219 B2RF	-	-	-	0.11	-	0.08	-	-	-	0.15	0.12	3.93	2.01	0.32	-	2.50	1.15	1.45
DP 1252 B2RF	0.08	-	0.41	-	-	4.07	-	-	-	-	0.45	-	-	3.94	-	0.17	-	0.70
DP 491	0.16	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*
Deltapine Total	**59.76**	**34.02**	**56.87**	**14.16**	**49.65**	**46.28**	**24.00**	**25.00**	**35.69**	**43.94**	**29.42**	**9.16**	**55.90**	**31.67**	**13.68**	**18.99**	**15.72**	**28.15**
Dyna-Gro																		
DG 2100 B2RF	-	0.03	-	-	-	-	-	-	-	-	-	-	-	-	-	0.23	-	0.13
DG 2400 RF	-	0.03	-	-	-	-	-	-	-	-	-	-	-	-	-	0.03	-	0.02
DG 2450 B2RF	-	-	-	-	-	-	-	-	-	-	0.39	-	0.46	-	-	0.12	-	0.09
DG 2520 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.34	-	0.19
DG 2570 B2RF	0.82	0.68	-	-	37.94	-	0.50	0.58	5.25	1.05	1.67	-	14.39	-	1.13	1.70	0.27	1.96
DG 2595 B2RF	-	-	-	-	-	-	-	-	-	-	-	-	0.46	-	-	-	-	0.01
Dyna-Gro Total	**0.82**	**0.74**	**-**	**-**	**37.94**	**-**	**0.50**	**0.58**	**5.25**	**1.05**	**2.05**	**-**	**15.31**	**-**	**1.13**	**2.43**	**0.27**	**2.39**
Phytogen																		
PHY 315 RF	0.03	0.56	-	-	-	-	-	0.42	-	-	0.03	-	-	0.02	-	-	0.10	0.04
PHY 367 WRF	0.68	-	3.04	-	-	0.08	-	-	1.81	-	2.49	13.35	0.44	1.70	2.75	2.27	9.60	1.71
PHY 375 WRF	7.96	8.14	4.12	-	-	5.73	4.50	1.35	9.56	3.69	13.54	18.90	0.37	12.20	39.11	2.36	24.86	5.44
PHY 440 W	0.42	-	-	-	-	1.06	-	-	-	-	-	-	-	-	-	0.38	-	0.34
PHY 485 WRF	-	0.64	-	-	-	-	-	-	-	1.49	-	-	-	-	-	0.01	-	0.10
PHY 499 WRF	17.42	3.70	3.07	-	8.62	32.39	-	19.70	-	3.85	28.07	28.76	0.22	37.55	6.37	3.40	24.76	9.65
PHY 565 WRF	5.00	-	0.94	-	-	2.06	-	-	-	-	0.20	-	-	4.33	-	*	-	0.52
PHY 72	-	-	-	3.82	-	-	-	-	-	-	-	-	-	-	-	-	-	0.05
PHY 725 RF	-	-	0.17	36.68	-	-	-	-	-	-	-	1.12	-	-	-	-	-	0.44
PHY 755 WRF	-	-	-	9.59	-	-	-	-	-	-	-	-	-	-	-	-	-	0.11
Phytogen Total	**31.51**	**13.05**	**11.34**	**50.09**	**8.62**	**41.33**	**4.50**	**21.48**	**11.37**	**9.03**	**44.32**	**62.13**	**1.03**	**55.80**	**48.24**	**8.42**	**59.32**	**18.40**
Seed Source Genetics																		
CT 110	-	-	-	-	-	-	-	-	0.04	-	-	-	-	-	-	-	-	*
Seed Source Genetics Total	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**0.04**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	**-**	*****
Miscellaneous																		
Misc.	0.83	-	4.33	3.52	-	0.20	3.50	-	0.10	-	0.13	4.51	0.71	0.38	0.16	0.31	0.18	0.40
Miscellaneous Total	**0.83**	**-**	**4.33**	**3.52**	**-**	**0.20**	**3.50**	**-**	**0.10**	**-**	**0.13**	**4.51**	**0.71**	**0.38**	**0.16**	**0.31**	**0.18**	**0.40**
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Transgenetic	**99.01**	**97.35**	**95.67**	**92.66**	**100.00**	**99.80**	**96.50**	**100.00**	**99.86**	**100.00**	**99.87**	**95.49**	**99.29**	**99.62**	**99.84**	**98.95**	**99.82**	**99.03**

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2012

Variety	US	Abilene				Corpus	Dumas			Florence			
	Total	KS	OK	TX	Classing Office	TX	AR	MS	Classing Office	NC	SC	VA	Classing Office
PHY 499 WRF Phytogen	9.65%	-	0.22%	2.19%	1.81%	9.08%	6.27%	3.26%	4.07%	28.07%	37.55%	24.76%	30.66%
DP 1044 B2RF Deltapine	7.16%	-	34.43%	21.51%	22.86%	14.92%	-	-	-	-	0.07%	-	0.02%
PHY 375 WRF Phytogen	5.44%	4.50%	0.37%	2.78%	2.45%	8.42%	1.87%	2.59%	2.40%	13.54%	12.20%	24.86%	14.18%
FM 9170 B2F Bayer CropScience - FiberMax	4.70%	-	0.32%	13.18%	10.76%	0.59%	-	-	-	-	-	-	-
DP 0912 B2RF Deltapine	4.28%	17.00%	4.05%	0.76%	1.78%	-	21.52%	6.44%	10.52%	3.08%	2.90%	1.46%	2.87%
FM 1740 B2F Bayer CropScience - FiberMax	4.11%	7.50%	15.42%	8.86%	9.85%	20.95%	-	-	-	0.18%	0.13%	0.53%	0.20%
ST 5458 B2RF Bayer CropScience - Stoneville	3.86%	-	-	2.30%	1.87%	1.44%	15.21%	22.70%	20.68%	0.30%	0.26%	-	0.26%
DP 1050 B2RF Deltapine	3.75%	-	-	-	-	1.36%	-	-	-	0.35%	4.99%	1.35%	1.87%
FM 2484 B2F Bayer CropScience - FiberMax	3.48%	-	0.33%	4.69%	3.86%	-	-	-	-	-	-	-	-
Epic RF All-Tex	3.17%	-	-	1.18%	0.96%	-	-	-	-	-	-	-	-
NG 4012 B2RF Americot	3.01%	-	0.61%	0.13%	0.20%	0.07%	-	-	-	-	-	-	-
DP 1048 B2RF Deltapine	2.92%	-	0.30%	0.14%	0.16%	2.83%	-	4.01%	2.93%	0.62%	5.94%	-	2.19%
ST 4145 LLB2 Bayer CropScience - Stoneville	2.74%	-	-	-	-	-	0.67%	5.32%	4.06%	13.68%	5.35%	10.22%	10.81%
ST 5288 B2F Bayer CropScience - Stoneville	2.11%	-	-	-	-	-	10.22%	13.29%	12.46%	4.03%	0.19%	3.79%	2.83%
DG 2570 B2RF Dyna-Gro	1.96%	0.50%	14.39%	3.72%	5.28%	0.24%	-	-	-	1.67%	-	0.27%	1.03%
FM 9063 B2F Bayer CropScience - FiberMax	1.94%	-	0.77%	8.17%	6.76%	-	-	-	-	-	-	-	-
NG 4111 RF Americot	1.77%	-	-	-	-	-	-	-	-	-	-	-	-
PHY 367 WRF Phytogen	1.71%	-	0.44%	1.05%	0.92%	1.29%	-	-	-	2.49%	1.70%	9.60%	2.91%
NG 3348 B2RF Americot	1.63%	2.50%	1.14%	0.74%	0.86%	-	-	-	-	-	-	-	-
FM 9180 B2F Bayer CropScience - FiberMax	1.45%	2.00%	0.44%	7.77%	6.45%	-	-	-	-	-	-	-	-
DP 1219 B2RF Deltapine	1.45%	-	2.01%	2.55%	2.38%	4.74%	-	-	-	0.12%	0.32%	1.15%	0.27%
DP 1137 B2RF Deltapine	1.43%	-	-	0.52%	0.42%	-	-	5.44%	3.97%	7.79%	8.98%	1.83%	7.60%
AM 1532 B2RF Americot	1.41%	-	2.03%	2.48%	2.33%	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2012

Variety	US	Lamesa	Lubbock	Macon				Memphis					
	Total	TX	TX	AL	FL	GA	Classing Office	AL	AR	MO	MS	TN	Classing Office
PHY 499 WRF Phytogen	9.65%	2.00%	2.46%	21.33%	8.62%	32.39%	29.46%	10.15%	2.44%	-	9.04%	6.37%	3.91%
DP 1044 B2RF Deltapine	7.16%	14.70%	5.26%	0.64%	-	-	0.09%	-	-	-	-	-	-
PHY 375 WRF Phytogen	5.44%	0.85%	0.46%	3.62%	-	5.73%	5.10%	16.02%	11.21%	9.56%	17.02%	39.11%	19.44%
FM 9170 B2F Bayer CropScience - FiberMax	4.70%	13.73%	8.10%	-	-	-	-	-	-	-	-	-	-
DP 0912 B2RF Deltapine	4.28%	0.91%	0.74%	0.64%	-	2.97%	2.48%	19.89%	26.88%	31.14%	20.48%	11.38%	22.70%
FM 1740 B2F Bayer CropScience - FiberMax	4.11%	1.17%	1.79%	1.29%	-	1.52%	1.39%	-	1.20%	0.86%	-	0.10%	0.62%
ST 5458 B2RF Bayer CropScience - Stoneville	3.86%	2.46%	1.59%	2.44%	-	1.83%	1.80%	0.55%	17.95%	25.23%	8.52%	0.19%	12.80%
DP 1050 B2RF Deltapine	3.75%	-	-	28.75%	26.21%	21.58%	22.82%	-	-	-	-	-	-
FM 2484 B2F Bayer CropScience - FiberMax	3.48%	9.73%	8.13%	-	-	-	-	-	-	-	-	-	-
Epic RF All-Tex	3.17%	1.44%	10.90%	-	-	-	-	-	-	-	-	-	-
NG 4012 B2RF Americot	3.01%	10.57%	8.30%	-	-	-	-	-	-	-	-	-	-
DP 1048 B2RF Deltapine	2.92%	-	-	24.04%	12.06%	13.16%	14.55%	-	0.21%	-	-	-	0.06%
ST 4145 LLB2 Bayer CropScience - Stoneville	2.74%	-	-	-	-	1.14%	0.92%	-	9.78%	12.13%	9.89%	29.47%	15.08%
ST 5288 B2F Bayer CropScience - Stoneville	2.11%	-	-	-	-	-	-	2.99%	2.36%	1.66%	21.43%	-	2.24%
DG 2570 B2RF Dyna-Gro	1.96%	1.39%	1.40%	1.26%	37.94%	-	2.48%	-	1.01%	5.25%	1.96%	1.13%	2.17%
FM 9063 B2F Bayer CropScience - FiberMax	1.94%	2.32%	3.00%	-	-	-	-	-	-	0.15%	-	0.04%	0.05%
NG 4111 RF Americot	1.77%	5.81%	5.04%	-	-	-	-	-	-	-	-	-	-
PHY 367 WRF Phytogen	1.71%	4.64%	2.39%	-	-	0.08%	0.07%	1.96%	-	1.81%	-	2.75%	1.50%
NG 3348 B2RF Americot	1.63%	7.16%	3.74%	-	-	-	-	-	-	-	-	-	-
FM 9180 B2F Bayer CropScience - FiberMax	1.45%	1.41%	1.58%	-	-	-	-	-	-	-	-	-	-
DP 1219 B2RF Deltapine	1.45%	1.99%	1.93%	-	-	0.08%	0.06%	-	-	-	1.81%	-	0.07%
DP 1137 B2RF Deltapine	1.43%	-	-	2.02%	3.79%	2.42%	2.45%	17.85%	-	-	5.10%	-	2.02%
AM 1532 B2RF Americot	1.41%	3.12%	3.20%	-	-	-	-	-	-	-	-	-	-

Table 2. Estimated percentage of upland cotton planted to specified varieties, by Classing Office and United States, crop of 2012

Variety	US	Rayville			Visalia				
	Total	LA	MS /1	Classing Office	AZ	CA	NM	TX /1	Classing Office
PHY 499 WRF Phytogen	9.65%	19.70%		18.67%	3.07%	-	28.76%		5.36%
DP 1044 B2RF Deltapine	7.16%	0.32%		0.41%	13.74%	-	5.22%		7.40%
PHY 375 WRF Phytogen	5.44%	1.35%		1.24%	4.12%	-	18.90%		6.00%
FM 9170 B2F Bayer CropScience - FiberMax	4.70%	-		-	0.45%	-	-		0.21%
DP 0912 B2RF Deltapine	4.28%	3.46%		4.02%	8.66%	-	-		4.12%
FM 1740 B2F Bayer CropScience - FiberMax	4.11%	-		-	6.20%	1.12%	-		3.35%
ST 5458 B2RF Bayer CropScience - Stoneville	3.86%	0.71%		0.65%	5.02%	-	-		2.39%
DP 1050 B2RF Deltapine	3.75%	0.51%		1.38%	2.84%	0.89%	-		1.67%
FM 2484 B2F Bayer CropScience - FiberMax	3.48%	-		-	0.04%	-	13.74%		2.14%
Epic RF All-Tex	3.17%	-		-	-	-	-		-
NG 4012 B2RF Americot	3.01%	-		-	-	-	-		-
DP 1048 B2RF Deltapine	2.92%	4.01%		4.16%	3.76%	0.56%	-		1.99%
ST 4145 LLB2 Bayer CropScience - Stoneville	2.74%	-		-	-	0.64%	-		0.23%
ST 5288 B2F Bayer CropScience - Stoneville	2.11%	51.24%		47.44%	1.66%	0.25%	-		0.88%
DG 2570 B2RF Dyna-Gro	1.96%	0.58%		2.58%	-	-	-		-
FM 9063 B2F Bayer CropScience - FiberMax	1.94%	-		-	1.14%	-	-		0.54%
NG 4111 RF Americot	1.77%	-		-	-	-	-		-
PHY 367 WRF Phytogen	1.71%	-		-	3.04%	-	13.35%		3.76%
NG 3348 B2RF Americot	1.63%	-		-	-	-	-		-
FM 9180 B2F Bayer CropScience - FiberMax	1.45%	-		-	1.07%	-	1.40%		0.68%
DP 1219 B2RF Deltapine	1.45%	-		*	-	0.11%	3.93%		0.51%
DP 1137 B2RF Deltapine	1.43%	0.99%		0.90%	-	-	-		-
AM 1532 B2RF Americot	1.41%	-		-	-	-	-		-

/1 Individual state data withheld

Table 3. Estimated percentage of American Pima cotton acreage planted to specified varieties, by states, crop of 2012

		AZ	CA	NM \1	TX \1	US
Phytogen	PHY 805 RF	42.31%	52.27%			48.80%
Phytogen	PHY 802 RF	9.23%	12.58%			12.68%
Deltapine	DP 340	21.76%	10.36%			12.25%
Phytogen	PHY 830	11.87%	8.38%			7.86%
Phytogen	PHY 800	11.87%	7.00%			6.60%
Deltapine	DP 357	-	3.55%			6.05%
Hazera	HA 195	-	3.28%			3.01%
Deltapine	DP 360	-	1.10%			1.00%
Hazera	HA 211	-	0.98%			0.90%
Miscellaneous	MISC	2.97%	0.30%			0.66%
Hazera	HA 222	-	0.20%			0.18%

/1 Individual state data withheld